

# ASTHMA FIRST AID

1



## SIT THE PERSON UPRIGHT

- Be calm and reassuring
- Do not leave them alone

2



## GIVE 4 SEPARATE PUFFS OF BLUE/GREY RELIEVER PUFFER

- Shake puffer
- Put 1 puff into spacer
- Take 4 breaths from spacer
  - Repeat until 4 puffs have been taken

OR give 2 separate inhalations of Bricanyl (6 years or older)

OR give 1 inhalation of Symbicort Turbuhaler (12 years or older)

OR give 2 puffs of Symbicort Rapihaler through a spacer (12 years or older)

**If no spacer available:** Take 1 puff as you take 1 slow, deep breath and hold breath for as long as comfortable. Repeat until all puffs are given

3



## WAIT 4 MINUTES

- If there is no improvement, give 4 more separate puffs of blue/grey reliever as above

OR give 1 more inhalation of Bricanyl

OR give 1 more inhalation of Symbicort Turbuhaler

OR give 2 puffs of Symbicort Rapihaler through a spacer

## IF THERE IS STILL NO IMPROVEMENT

4



## DIAL TRIPLE ZERO (000)

- Say 'ambulance' and that someone is having an asthma attack
- Keep giving 4 separate puffs every 4 minutes until emergency assistance arrives

OR give 1 inhalation of a Bricanyl or Symbicort Turbuhaler every 4 minutes – up to a max of 4 more inhalations of Symbicort Turbuhaler

OR give 2 puffs of Symbicort Rapihaler through a spacer every 4 minutes – up to a max of 8 more puffs of Symbicort Rapihaler

## CALL EMERGENCY ASSISTANCE IMMEDIATELY AND DIAL TRIPLE ZERO (000) IF:

- the person is not breathing
- the person's asthma suddenly becomes worse or is not improving
- the person is having an asthma attack and a reliever is not available
- you are not sure if it is asthma
- **the person is known to have anaphylaxis – follow their Anaphylaxis Action Plan, then give Asthma First Aid**

Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma.



Translating and Interpreting Service  
131 450



**ASTHMA AUSTRALIA**

**1800 ASTHMA**  
(1800 278 462)

[asthma.org.au](http://asthma.org.au)