# <u>ASTHMA FIRST AID</u>





### SIT THE PERSON UPRIGHT

- Be calm and reassuring
- Do not leave them alone

# 2



### GIVE 4 SEPARATE PUFFS OF BLUE/GREY RELIEVER PUFFER

- Shake puffer
- Put 1 puff into spacer
- Take 4 breaths from spacer
  - Repeat until 4 puffs have been taken
- OR give 2 separate inhalations of Bricanyl (6 years or older)
- OR give 1 inhalation of Symbicort Turbuhaler (12 years or older)
- OR give 2 puffs of Symbicort Rapihaler through a spacer (12 years or older)

If no spacer available: Take 1 puff as you take 1 slow, deep breath and hold breath for as long as comfortable. Repeat until all puffs are given

# 3



#### **WAIT 4 MINUTES**

- If there is no improvement, give 4 more separate puffs of blue/grey reliever as above
- OR give 1 more inhalation of Bricanyl
- OR give 1 more inhalation of Symbicort Turbuhaler
- OR give 2 puffs of Symbicort Rapihaler through a spacer

### IF THERE IS STILL NO IMPROVEMENT





### **DIAL TRIPLE ZERO (000)**

- Say 'ambulance' and that someone is having an asthma attack
- Keep giving 4 separate puffs every 4 minutes until emergency assistance arrives

OR give 1 inhalation of a Bricanyl or Symbicort Turbuhaler every 4 minutes

– up to a max of 4 more inhalations of Symbicort Turbuhaler

OR give 2 puffs of Symbicort Rapihaler through a spacer every 4 minutes

– up to a max of 8 more puffs of Symbicort Rapihaler

## CALL EMERGENCY ASSISTANCE IMMEDIATELY AND DIAL TRIPLE ZERO (000) IF:

- the person is not breathing
- the person's asthma suddenly becomes worse or is not improving
- the person is having an asthma attack and a reliever is not available
- you are not sure if it is asthma
- the person is known to have anaphylaxis follow their Anaphylaxis Action Plan, then give Asthma First Aid

Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma.



Translating and Interpreting Service 131 450



1800 ASTHMA

(1800 278 462)

asthma.org.au