

LIVE WELL WITH YOUR ASTHMA



COULD IT BE ASTHMA?

Do you ever:

- Get breathless or wheezy?
- Wake up coughing or wheezing?
- Struggle to keep up with your normal activity?

Being breathless is not a normal part of life or getting older. Ask your pharmacist or doctor about a lung health check.

IF YOU HAVE ASTHMA, SEE YOUR DOCTOR IF YOU:

- Are coughing, wheezing or breathless at any time of the day or night
- Struggle to keep up with normal activity
- Use your blue/grey reliever puffer more than two days per week
- Are unsure about the way your medications and devices can work best for you
- Have had any asthma symptoms in the past month or a flare-up in the past year and you are not currently on asthma preventer medication

ASTHMA SHOULD NOT STOP YOU LIVING YOUR LIFE WELL!



All Asthma Australia information is consistent with the National Asthma Council's, The Australian Asthma Handbook (2020).

Asthma Australia information does not replace professional medical advice. People should ask their doctor any questions about diagnosis and treatment.

©Asthma Australia 2022.

This work is licensed under the Creative Commons Attribution-NonCommercial 3.0 Australia License.



ASTHMA FIRST AID

Blue/Grey Reliever

Airomir, Asmol, Ventolin or Zempreon and Bricanyl

Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma



DIAL TRIPLE ZERO (000) FOR AN AMBULANCE IMMEDIATELY IF THE PERSON:

- is not breathing
- suddenly becomes worse or is not improving
- is having an asthma attack and a reliever is not available
- is unsure if it is asthma
- **has a known allergy to food, insects or medication and has SUDDEN BREATHING DIFFICULTY, GIVE ADRENALINE AUTOINJECTOR FIRST (if available), even if there are no skin changes, then use a reliever**

1



SIT THE PERSON UPRIGHT

- Be **calm** and reassuring
- **Do not leave** them alone

2



GIVE 4 SEPARATE PUFFS OF RELIEVER PUFFER

- **Shake** puffer
- Put **1 puff** into spacer
- Take **4 breaths** from spacer
 - Repeat until **4 puffs** have been taken



If using **Bricanyl**, give 2 separate inhalations (5 years or older)

If you don't have a spacer handy in an emergency, take **1 puff** as you take **1 slow, deep breath** and hold breath for as long as comfortable. **Repeat** until all puffs are given

3



WAIT 4 MINUTES

- If breathing does not return to normal, give **4 more separate puffs** of reliever as above



Bricanyl: Give 1 more inhalation

IF BREATHING DOES NOT RETURN TO NORMAL

4



DIAL TRIPLE ZERO (000)

- Say **'ambulance'** and that someone is having an asthma attack
- Keep giving **4 separate puffs every 4 minutes** until emergency assistance arrives



Bricanyl: Give 1 more inhalation **every 4 minutes** until emergency assistance arrives