ASTHMA CARE PLAN FOR EDUCATION AND CARE SERVICES

CONFIDENTIAL: Staff are trained in Asthma First Aid (see overleaf) and can provide routine asthma medication as authorised in this care plan by the treating doctor. Please advise staff in writing of any changes to this plan.

To be completed by the treating doctor and parent/guardian, for supervising staff and emergency medical personnel.

and emergency medical personnel.

PLEASE PRINT CLEARLY

Student's name: ______ DOB: ______

PHOTO OF STUDENT (OPTIONAL)

Plan date /20	
Review date	

MANAGING AN ASTHMA ATTACK

Staff are trained in Asthma First Aid (see overleaf). Please write down anything different this student might need if they have an asthma attack:

DAILY ASTHMA MANAGEMENT

This student's usual ast	nma signs:	Frequency and severity:		Known triggers for this student's asthma (e.g. exercise*, colds/flu, smoke) —
Cough		Daily/most days		please detail:
Wheeze		Frequently (more than 5 x per	year)	
Difficulty breathing	5	Occasionally (less than 5 x pe	r year)	
Other (please describe):		Other (please describe)		
Does this student usually	/ tell an adult if s,	he is having trouble breathing?	Yes	No
Does this student need help to take asthma medication?			Yes	No
Does this student use a	mask with a spac	er?	Yes	No
*Does this student need a blue/grey reliever puffer medication before exercise?			Yes	No

MEDICATION PLAN

Signature

NAME OF MEDICATION AND COLOUR

If this student needs asthma medication, please detail below and make sure the medication and spacer/mask are supplied to staff.

DOSE/NUMBER OF PUFFS

DOCTOR	PARENT/GUARDI	EMERGENCY CONTACT INFORMATION			
Name of doctor	attachments listed. I appro	d agreed with this care plan and any ve the release of this information to al personnel. I will notify the staff in	Contact name		
Address	writing if there are any chang staff will seek emergency I am responsible for paymer	Phone			
Phone	Signature	Date	Mobile		

ASTHMA AUSTRALIA

Email

TIME REQUIRED

Name

Date

ASTHMA FIRST AID

Blue/Grey Reliever

Airomir, Asmol, Ventolin or Zempreon and Bricanyl

Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma



DIAL TRIPLE ZERO (000) FOR AN AMBULANCE IMMEDIATELY IF THE PERSON:

- is not breathing
- suddenly becomes worse or is not improving
- is having an asthma attack and a reliever is not available
- is unsure if it is asthma
- has a known allergy to food, insects or medication and has SUDDEN BREATHING DIFFICULTY, GIVE ADRENALINE AUTOINJECTOR FIRST (if available)





SIT THE PERSON UPRIGHT

- Be calm and reassuring
- Do not leave them alone

2



GIVE 4 SEPARATE PUFFS OF RELIEVER PUFFER

- Shake puffer
- Put 1 puff into spacer
- Take 4 breaths from spacer
 - Repeat until 4 separate puffs have been taken



If using **Bricanyl** (5 years or older)

- Do not shake. Open, twist around and back, and take a deep breath in
- Repeat until 2 separate inhalations have been taken

If you don't have a spacer handy in an emergency, take <u>1 puff</u> as you take <u>1 slow, deep breath</u> and hold breath for as long as comfortable. **Repeat** until all puffs are given

3



WAIT 4
MINUTES

If breathing does not return to normal, give
 4 more separate puffs of reliever as above



Bricanyl: Give 1 more inhalation

IF BREATHING DOES NOT RETURN TO NORMAL





DIAL TRIPLE ZERO (000)

- Say 'ambulance' and that someone is having an asthma attack
- Keep giving 4 separate puffs every
 4 minutes until emergency assistance arrives



Bricanyl: Give 1 more inhalation <u>every 4 minutes</u> until emergency assistance arrives







