

ASTHMA ACTION PLAN




VICTORIAN SCHOOLS

Student's name: _____

DOB: _____

Confirmed triggers: _____



- Child can self-administer if well enough
- Child needs to pre-medicate prior to exercise
- Face mask needed with spacer

ALWAYS give adrenaline autoinjector FIRST, and then asthma reliever puffer if someone with known asthma and allergy to food, insects or medication has SUDDEN BREATHING DIFFICULTY (including wheeze, persistent cough or hoarse voice) even if there are no skin symptoms.

Adrenaline autoinjector prescribed: Y N Type of adrenaline autoinjector: _____

ASTHMA FIRST AID

For Severe or Life-Threatening signs and symptoms, call for emergency assistance immediately on Triple Zero "000"
Mild to moderate symptoms do not always present before severe or life-threatening symptoms

- 1. Sit the person upright**
Stay with the person and be calm and reassuring
- 2. Give 2 separate puffs of Symbicort**
Shake the puffer before each puff
Puff 1 puff into the spacer at a time
Take 4 breaths from spacer between each puff
- 3. Wait 4 minutes**
If there is no improvement, give 2 doses of Symbicort
- 4. If there is still no improvement call emergency assistance**
Dial Triple Zero "000"
Say 'ambulance' and that someone is having an asthma attack
Keep giving 2 puffs every 4 minutes until emergency assistance arrives (**maximum 12 doses total**)

If maximum dose is reached before emergency services arrive follow the 4 x 4 Asthma First Aid plan on reverse.

Commence CPR at any time if person is unresponsive and not breathing normally.

SIGNS AND SYMPTOMS

MILD TO MODERATE

- Minor difficulty breathing
- May have a cough
- May have a wheeze
- Other signs to look for:



SEVERE

- Cannot speak a full sentence
- Sitting hunched forward
- Tugging in of skin over chest/throat
- May have a cough or wheeze
- Obvious difficulty breathing
- Lethargic
- Sore tummy (young children)

LIFE-THREATENING

- Unable to speak or 1-2 words
- Collapsed/exhausted
- Gasping for breath
- May no longer have a cough or wheeze
- Drowsy/confused/unconscious
- Skin discolouration (blue lips)

Emergency contact name:

Work ph:

Home ph:

Mobile ph:

Plan prepared by Dr or Nurse Practitioner:

Signed: I hereby authorise medications specified on this plan to be administered according to the plan

Date prepared:

Date of next review:



- Assemble spacer.
- Remove cap from puffer.
- Shake puffer well.
- Attach puffer to end of spacer.
- Place mouthpiece of spacer in mouth and ensure lips seal around it.
- Breathe out gently into the spacer.
- Press down on puffer canister once to fire medication into spacer.
- Breathe in and out normally for 4 breaths (keeping your mouth on the spacer).

ASTHMA FIRST AID

Dual Purpose Reliever Symbicort Rapihaler 100/3



DIAL TRIPLE ZERO (000) FOR AN AMBULANCE IMMEDIATELY IF THE PERSON:

- is not breathing
- suddenly becomes worse or is not improving
- is having an asthma attack and a reliever is not available
- is unsure if it is asthma
- has a known allergy to food, insects or medication and has **SUDDEN BREATHING DIFFICULTY**, GIVE ADRENALINE AUTOINJECTOR FIRST (if available), even if there are no skin changes, then use a reliever

1



SIT THE PERSON UPRIGHT

- Be calm and reassuring
- Do not leave them alone

2



GIVE 2 PUFFS OF SYMBICORT

- Shake puffer
- Put 1 puff into spacer
- Take 4 breaths from spacer
 - Repeat until 2 puffs have been taken

If you don't have a spacer handy in an emergency, take 1 puff as you take 1 slow, deep breath and hold breath for as long as comfortable. **Repeat** until all puffs are given

3



WAIT 4 MINUTES

- If breathing does not return to normal, give 2 more puffs of **Symbicort** through a spacer

IF BREATHING DOES NOT RETURN TO NORMAL

4



DIAL TRIPLE ZERO (000)

- Say 'ambulance' and that someone is having an asthma attack
- Give 2 more puffs of **Symbicort** through a spacer every 4 minutes until emergency assistance arrives
 - up to a maximum of 8 more puffs