

# ASTHMA FIRST AID

## Blue/Grey Reliever

Airomir, Asmol, Ventolin or Zempreon and Bricanyl

Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma



**DIAL TRIPLE ZERO (000) FOR AN AMBULANCE IMMEDIATELY IF THE PERSON:**

- is not breathing
- suddenly becomes worse or is not improving
- is having an asthma attack and a reliever is not available
- is unsure if it is asthma
- **has a known allergy to food, insects or medication and has SUDDEN BREATHING DIFFICULTY, GIVE ADRENALINE AUTOINJECTOR FIRST (if available)**

**1**



**SIT THE PERSON UPRIGHT**

- Be calm and reassuring
- Do not leave them alone

**2**



**GIVE 4 SEPARATE PUFFS OF RELIEVER PUFFER**

- Shake puffer
- Put 1 puff into spacer
- Take 4 breaths from spacer
  - Repeat until 4 separate puffs have been taken



If using **Bricanyl** (5 years or older)

- **Do not shake.** Open, twist around and back, and take a deep breath in
- Repeat until 2 separate inhalations have been taken

If you don't have a spacer handy in an emergency, take 1 puff as you take 1 slow, deep breath and hold breath for as long as comfortable. Repeat until all puffs are given

**3**



**WAIT 4 MINUTES**

- If breathing does not return to normal, give 4 more separate puffs of reliever as above



**Bricanyl:** Give 1 more inhalation

## IF BREATHING DOES NOT RETURN TO NORMAL

**4**



**DIAL TRIPLE ZERO (000)**

- Say 'ambulance' and that someone is having an asthma attack
- Keep giving 4 separate puffs every 4 minutes until emergency assistance arrives



**Bricanyl:** Give 1 more inhalation every 4 minutes until emergency assistance arrives



**ASTHMA AUSTRALIA**

**1800 ASTHMA**  
(1800 278 462)  
[asthma.org.au](http://asthma.org.au)

Supported by:



Translating and Interpreting Service  
131 450